

Award Number: W81XWH-16-1-0239

**TITLE:**

**Lung Cancer-Specific Circular RNAs as Biomarkers**

**PRINCIPAL INVESTIGATOR:**

Yin-Yuan Mo

**CONTRACTING ORGANIZATION:**

University of Mississippi Medical Center  
Jackson, MS 39216

**REPORT DATE:**

August 2017

**TYPE OF REPORT:**

Annual

**PREPARED FOR:** U.S. Army Medical Research and Materiel Command  
Fort Detrick, Maryland 21702-5012

**DISTRIBUTION STATEMENT:** Approved for Public Release;  
Distribution Unlimited

The views, opinions and/or findings contained in this report are those of the author(s) and should not be construed as an official Department of the Army position, policy or decision unless so designated by other documentation.

REPORT DOCUMENTATION PAGE				Form Approved OMB No. 0704-0188	
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing this collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number. <b>PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.</b>					
1. REPORT DATE August 2017		2. REPORT TYPE Annual		3. DATES COVERED 1 Aug 2016 - 31 Jul 2017	
4. TITLE AND SUBTITLE  Lung Cancer-Specific Circular RNAs as Biomarkers				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER W81XWH-16-1-0239	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Yin-Yuan Mo  E-Mail: ymo@umc.edu				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES)  University of Mississippi Medical Center 2500 State Street, G652 Jackson, MS 39216				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES) U.S. Army Medical Research and Materiel Command Fort Detrick, Maryland 21702-5012				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION / AVAILABILITY STATEMENT Approved for Public Release; Distribution Unlimited					
13. SUPPLEMENTARY NOTES					
14. ABSTRACT  The major goal of this application is to determine whether lung cancer cells differentially express circular RNAs such that these circular RNAs may serve as novel biomarkers for lung cancer diagnosis and prognosis. There are two Tasks in this application. First, we will perform microarray profiling to determine whether lung cancer cells display different patterns of circular RNAs from those of normal cells. Second, we will determine whether differential expression of circular RNAs can also be detected in cell culture models. Third, we will determine whether circular RNAs can be detected in circulation system. The success of this study may lead to novel clinical applications such as identification of biomarkers or therapeutic targets for lung cancer.					
15. SUBJECT TERMS lncRNA, circRNA, biomarker					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT	18. NUMBER OF PAGES	19a. NAME OF RESPONSIBLE PERSON
a. REPORT	b. ABSTRACT	c. THIS PAGE			USAMRMC
U	U	U	UU	7	19b. TELEPHONE NUMBER (include area code)

## Table of Contents

	<u>Page</u>
Introduction.....	1
Body.....	1
Key Research Accomplishments.....	3
Reportable Outcomes.....	N/a
Conclusion.....	4
References.....	N/a
Appendices.....	N/a

## Introduction

Circular RNAs (circRNAs) belong to a special group of long non-coding RNAs (lncRNAs). Although in some cases, circRNAs can code for protein, most of them lack the coding capacity. circRNAs are formed from lncRNAs or protein coding genes often through back splicing. There are five types of circRNA derived from: 1) exonic, 2) intronic, 3) antisense, 4) sense overlapping and 5) intergenic. "Exonic" represents circRNA arising from the exons of the linear transcript; "Intronic" represents the circRNA arising from an intron of the linear transcript; "antisense" represents circRNA whose gene locus overlap with the linear RNA, but transcribed from the opposite strand; "sense overlapping" represents circRNA transcribed from same gene locus as the linear transcript, but not classified into "exonic" and "intronic"; lastly, "intergenic" represents circRNA located outside known gene locus.

Despite different types of circRNAs, they all can play a regulatory role in gene expression through microRNA-mediated repression. Thus, we hypothesize that lung cancer may exploit this mechanism for its own advantage and as such lung cancer may display a very different circRNA pattern from normal lung cells. Therefore, the major goal of this application is to determine whether we can identify differentially expressed circRNAs in lung cancer.

## Body

**CircRNAs are aberrantly expressed in lung cancer.** As newly discovered molecules, circRNAs are poorly characterized. Little is known whether they are dysregulated in lung cancer. Thus, our first step was to characterize these new molecules by profiling. Results

indicate that over one hundred of circRNAs are either upregulated or downregulated. We listed four of them as representative differentially expressed circRNAs in Table 1.

Table 1, Representative circRNAs that are differentially expressed (tumor cells vs normal)

circRNA	Fold change	Regulation	Transcript	MRE1	MRE2	MRE3	MRE4	MRE5
hsa_circRNA_400633	8.8	up	NM_005063	<a href="#">hsa-miR-5589-5p</a>	<a href="#">hsa-miR-1253</a>	<a href="#">hsa-miR-3127-3p</a>	<a href="#">hsa-miR-541-3p</a>	<a href="#">hsa-miR-3691-3p</a>
hsa_circRNA_101100	8.6	up	NM_007007	<a href="#">hsa-miR-183-5p</a>	<a href="#">hsa-miR-383-3p</a>	<a href="#">hsa-miR-433-3p</a>	<a href="#">hsa-miR-607</a>	<a href="#">hsa-miR-640</a>
hsa_circRNA_005054	0.02	down	NM_001103184	<a href="#">hsa-miR-612</a>	<a href="#">hsa-miR-6515-3p</a>	<a href="#">hsa-miR-4753-5p</a>	<a href="#">hsa-miR-362-5p</a>	<a href="#">hsa-miR-450a-2-3p</a>
hsa_circRNA_001831	0.03	down	NM_015356	<a href="#">hsa-miR-3151-5p</a>	<a href="#">hsa-miR-6791-5p</a>	<a href="#">hsa-miR-939-5p</a>	<a href="#">hsa-miR-637</a>	<a href="#">hsa-miR-7974</a>

For example, hsa\_circRNA\_400633 and hsa\_circRNA\_101100 were upregulated with over an 8-fold increase in tumor vs normal with p value <0.05. On the other hand, the expression level of hsa\_circRNA\_005054 and hsa\_circRNA\_001831 was 0.02 and

0.03, respectively, as  
 compared to normal  
 cells as 1. To better  
 illustrate how the  
 circular form is  
 formed, we provide the  
 sequence for  
 hsa\_circRNA\_400633  
 and  
 hsa\_circRNA\_101100,




Fig. 1 DNA sequence of hsa\_circRNA\_400633. The junction of two ends is highlighted by red and blue, respectively.

as shown in Figs. 1 and 2 as an example. The top part is the actual sequence and the bottom part is when a circle is formed. Two ends at the junction were highlighted by green and blue, respectively.

**CircRNAs can potentially target microRNAs.** One of potential functions for circRNAs is the capability to serve as sponges to neutralize the endogenous microRNAs. In this

gaagggaaatggatactgcaagaacgccattgagtgaagctgaatttgaagaaatcatgaataga  
aatagggcaatctcaagcagtgctatcttcgagagctgtgtctgatgccagtgctggtgattatgg  
gagtgtctattgagacactggtaactgcaatttctttaattaacaatccaaagtatctgctgatg  
atcgttgcaaagttcttattagttctttgcaagattgccttcattggaattgagtccaagtcttat  
ggttctggatcaagacgtgaacgatcaagagagagggaccatagtagatcacgagaaaagagtcg  
acgtcataaatcccgtagtagagaccgtcatgacgattattacagagagagagaagcagagaaacgag  
agaggcaccgggatcgtgaccgagaccgtgaccgagagcgtgaccgagagcgcgaatatcgatcat  
cgttta




Fig. 2 DNA sequence of hsa\_circRNA\_101100. The junction of two ends is highlighted by blue and green, respectively.

microRNAs each for four representative circRNAs (Table 1). For example, miR-541-3p suppresses tumor progression by directly targeting TGIF2 in non-small cell lung cancer. On the other hand, miR-183 regulates autophagy and apoptosis in colorectal cancer through targeting of UVRAG; and it promotes proliferation and invasion in esophageal squamous cell carcinoma by targeting programmed cell death 4. It is evident that these microRNAs can serve as either oncogenes or tumor suppressors, thus impacting various aspect of tumorigenesis.

Together, these findings suggest that aberrant expression of these circRNAs may affect the levels of these microRNAs, thus, contributing to lung tumorigenesis.

## Key Research Accomplishments

regard, all  
of these  
circRNAs  
had the  
potential to  
target more  
than one  
microRNAs.  
We listed  
five

- We identified over 100 upregulated or downregulated circRNAs from lung cancer cells through profiling.
- All of these identified circRNAs carry microRNA binding sites, through which they may regulate the level of endogenous microRNAs.
- In this regard, hsa\_circRNA\_400633 can potentially target miR-541-3p which has been shown to be a tumor suppressive microRNA.
- We will determine whether any of these circRNAs impact tumor cell growth in the cell culture models. We will also test whether they are differentially expressed in serum samples from normal and lung cancer patients such that they may serve as biomarkers for lung cancer.

### **Reportable Outcomes**

Not yet.

### **Conclusions**

Microarray profiling has identified over a hundred of upregulated or downregulated circRNAs from lung cancer cells. We are currently determining whether ectopic expression of any of these circRNAs will impact lung tumor cell growth and invasion. We will also determine these circRNAs can target the predicted microRNAs which in turns regulate their expression. Finally, we will test their potential as lung cancer biomarkers.